

SUPER-GEOMETRIC CONVERGENCE OF A SPECTRAL ELEMENT METHOD FOR EIGENVALUE PROBLEMS WITH JUMP COEFFICIENTS*

Lin Wang

*College of Mathematics and Computer Science, Hunan Normal University, Changsha 410081, China
and College of Information Technical Science, Nankai University,
Tianjin 300071, China*

Email: rinoya@mail.nankai.edu.cn

Ziqing Xie

*College of Mathematics and Computer Science, Hunan Normal University,
Changsha 410081, China*

Email: ziqingxie@hunnu.edu.cn

Zhimin Zhang

*Department of Mathematics, Wayne State University, Detroit, MI 48202, USA
and College of Mathematics and Computer Science, Hunan Normal University,
Changsha 410081, China*

Email: zzhang@math.wayne.edu

Abstract

We propose and analyze a C^0 spectral element method for a model eigenvalue problem with discontinuous coefficients in the one dimensional setting. A super-geometric rate of convergence is proved for the piecewise constant coefficients case and verified by numerical tests. Furthermore, the asymptotical equivalence between a Gauss-Lobatto collocation method and a spectral Galerkin method is established for a simplified model.

Mathematics subject classification: Primary 65N30, Secondary 65N50, 65N15, 65N12, 65D10, 74S05, 41A10, 41A25.

Key words: Eigenvalue, Spectral method, Collocation, Galerkin finite element method.

1. Introduction

We often encounter eigenvalue problems with discontinuous coefficients in practice. Examples of such applications may be found in [11]. In this paper, we consider the following one dimensional model problem: Find $(\lambda, u) \in \mathbb{R}^+ \times H^2(-\pi, \pi)$ such that

$$-u''(x) = \lambda c(x)u(x), \quad u(-\pi) = u(\pi), \quad u'(-\pi) = u'(\pi). \quad (1.1)$$

Here $c(x) \geq c_0 > 0$ is a piecewise constant, or piecewise analytic function. The physics background of this model problem comes from the source-free Maxwell equations describing the transverse-magnetic mode in the one-dimensional periodic media, where the function u represents the electric field pattern, and the dielectric function $c(x)$ describes a unit cell from a multilayer structure with 2π -periodicity. This model problem was discussed by Min and

* Received March 18, 2009 / Revised version received May 5, 2009 / Accepted June 6, 2009 /
Published online February 1, 2010 /

Gottlieb in [11] where C^1 conforming spectral collocation methods were constructed on two elements over

$$H_{per}^2(-\pi, \pi) = \{v \in H^2(-\pi, \pi) : v(-\pi) = v(\pi), v'(-\pi) = v'(\pi)\},$$

and error bounds of type $\mathcal{O}(p^{-m})$ were established. Note that the solution of (1.1) belongs to C^1 .

It would be interesting to discuss C^0 spectral element methods over

$$H_{per}^1(-\pi, \pi) = \{v \in H^1(-\pi, \pi) : v(-\pi) = v(\pi)big\},$$

since the construction of a C^0 spectral element method is much simpler than that of the global C^1 spectral collocation method proposed in [11]. The idea of the spectral element can be found, e.g., in an early work [12]. Note that the spectral element method is equivalent to the so-called p -version finite element method, see e.g., [3]. Under the finite element variational framework, we are able to prove a super-geometric error bound of type $\mathcal{O}(e^{-2p(\log p-\gamma)})$. In some earlier works of the third author, the super-geometric error bound of type $\mathcal{O}(e^{-p(\log p-\gamma)})$ has been established for some spectral/collocation approximations of the two-point boundary problem [17, 18]. Our error bound for the eigenvalue approximation “doubles” the error bound for the associated eigenfunction approximation, the fact we have known for the h -version finite element method. It is worthy to point out that in the literature of the spectral method, it is a common practice to consider error bounds of type $\mathcal{O}(p^{-m})$, see, e.g., [5–7, 10, 15, 16], and reference therein. To the best of our knowledge, this is the first time that a super-geometric convergence rate is established for the eigenvalue approximation by the spectral method.

2. Theoretical Setting

The variational formulation of (1.1) is to find $(\lambda, u) \in \mathbb{R}^+ \times H_{per}^1(-\pi, \pi)$ such that

$$(u', v') = \lambda(cu, v), \quad \forall v \in H_{per}^1(-\pi, \pi). \tag{2.1}$$

In this paper, we also consider the Dirichlet problem

$$-u''(x) = \lambda c(x)u(x), \quad u(0) = 0 = u(1).$$

Its variational formulation is to find $(\lambda, u) \in \mathbb{R}^+ \times H_0^1(0, 1)$ such that

$$(u', v') = \lambda(cu, v), \quad \forall v \in H_0^1(0, 1). \tag{2.2}$$

By the general theory [2, 8], both problems (2.1) and (2.2) have countable infinite sequence of eigen-pairs (λ_j, u_j) satisfying

$$0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots \rightarrow \infty, \quad (u'_i, u'_j) = \lambda_j(cu_i, u_j) = \lambda_j \delta_{ij}.$$

Furthermore, eigenvalues can be characterized as extrema of the Rayleigh quotient $R(u) = (u', u')/(cu, u)$ as follows

$$\lambda_1 = \inf_{u \in S} R(u),$$

$$\lambda_k = \inf_{u \in S, (u', u'_j)=0, j=1, \dots, k-1} R(u) = R(u_k), \quad k = 2, 3, \dots,$$

where $S = H_{per}^1(-\pi, \pi)$ or $H_0^1(0, 1)$.

Next, we describe the framework of our numerical approximation. We partition the solution interval into m sub-intervals (element) such that $c(x)$ is analytic on each interval. Let h be the maximum length of all elements, we then define a finite dimensional subspace $S_p^h \subset S$, as a piecewise polynomial of degree p on each element. Our spectral element method is to find an eigen-pair $(\lambda(p), w_p) \in R^+ \times S_p^h$ such that

$$(w'_p, v') = \lambda(p)(cw_p, v), \quad \forall v \in S_p^h. \tag{2.3}$$

Note that the partition parameter h is fixed and convergence is achieved by increasing polynomial degree p . Therefore, we may suppress the index h later.

By the general theory [2, 8], the problem (2.3) has a finite sequence of eigen-pairs $(\lambda_{j,p}, w_{j,p})$ satisfying

$$\begin{aligned} 0 < \lambda_{1,p} \leq \lambda_{2,p} \leq \dots \leq \lambda_{N,p}, \quad N = \begin{cases} mp - 1 & H_0^1(0, 1) \\ mp & H_{per}^1(-\pi, \pi) \end{cases} \\ (w'_{i,p} w'_{j,p}) = \lambda_{j,p}(cw_{i,p}, w_{j,p}) = \lambda_{j,p} \delta_{ij}; \\ \lambda_{1,p} = \min_{w \in S_p} = R(w_{1,p}), \end{aligned} \tag{2.4}$$

$$\lambda_{k,p} = \min_{w \in S_p, (w', w'_{j,p})=0, j=1, \dots, k-1} R(w) = R(w_{k,p}), \quad k = 2, 3, \dots \tag{2.5}$$

One important observation from the above Minimum-Maximum principle is that the specific eigenvalue approximation is from above in the sense

$$\lambda_k \leq \dots \leq \lambda_{k,p+1} \leq \lambda_{k,p} \leq \lambda_{k,p-1} \dots \leq \lambda_{k,1}.$$

3. A Galerkin Spectral (p-Version) Method

Without loss of generality, we consider piecewise constant $c(x)$ as in [11] with jump at the center of the solution domain. In particular, we take

$$c(x) = \begin{cases} 1 & x \in (-\pi, 0), \\ \omega^2 & x \in (0, \pi). \end{cases}$$

Instead of constructing C^1 shape functions for eigenvalue problem (2.1), we seek for a C^0 approximation $w_p \in H^1(0, 1)$ with traditional expansion

$$w_p(x) = w^0(N_- + N_+) + \sum_{j=1}^{p-1} w^j \phi_{p-j+1}(x) + w^p N(x) + \sum_{j=p+1}^{2p-1} w^j \psi_{j-p+1}(x), \tag{3.1}$$

where $N_-(x), N(x)$, and $N_+(x)$ are linear nodal shape functions at the left end, middle point, and right end of the solution interval, respectively; ϕ_j and ψ_j are bubble functions on the left and right intervals, respectively. The counterpart of ϕ_{k+1} in $[-1, 1]$ is defined as

$$\hat{\phi}_{k+1}(\xi) = \sqrt{\frac{2k+1}{2}} \int_{-1}^{\xi} L_k(t) dt = \frac{1}{\sqrt{2(2k+1)}} \left(L_{k+1}(\xi) - L_{k-1}(\xi) \right). \tag{3.2}$$

and the counterpart of ψ_{k+1} in $[-1, 1]$ is defined similarly. Note that $w^0 = 0$ for the eigenvalue problem (2.2). With this setting, the resulting stiffness matrix is diagonal and the mass matrix is 5-diagonal, see Appendix.

4. Super-Geometric Convergence Rate

Let (λ_k, u_k) be the k th eigen-pair and $(\lambda_{k,h}, u_{k,h}) \in R \times S^h$ be its h -version finite element approximation. According to [2, p.700],

$$C_1 \epsilon_h^2 \leq \lambda_{k,h} - \lambda_k \leq C_2 \epsilon_h^2,$$

with

$$\epsilon_h = \inf_{\chi \in S^h} \|u_k - \chi\|_1,$$

for simple eigenvalue λ_k (see [2, p.695 (8.21)]). Transferring this theory to our spectral element method language, we have, for any simple eigen-pair (λ, u) ,

$$C_1 \epsilon_p^2 \leq \lambda_p - \lambda \leq C_2 \epsilon_p^2, \tag{4.1}$$

with

$$\epsilon_p = \inf_{\chi \in S_p} \|u - \chi\|_1 \approx \inf_{\chi \in S_p} \|u' - \chi'\| \approx \|u' - u'_p\|,$$

where $u_p \in S_p$ such that u'_p is the piecewise Legendre expansion of u' (not solution of (2.1) or (2.2)). Note that the first “ \approx ” comes from the Poincaré inequality and the last “ \approx ” is based on the fact that the Legendre expansion minimizes the L^2 -norm.

Lemma 4.1. *Let u satisfy the regularity assumption*

$$\max_{x \in [-1,1]} |u^{(k)}(x)| \leq cM^k$$

for fixed constants c and M , and let \tilde{u}'_p be the Legendre expansion of u' on $[-1, 1]$. Then under the assumption $(2p + 1)(2p + 3) > 2M^2$,

$$\|u' - \tilde{u}'_p\|_{L_2[-1,1]} \leq C\sqrt{p} \left(\frac{eM}{2p}\right)^{p+1}, \tag{4.2}$$

where C is independent of p and M .

Proof. The error of $(p - 1)$ -term Legendre expansion is

$$\|u' - \tilde{u}'_p\|_{L_2[-1,1]}^2 = \sum_{k=p}^{\infty} \frac{2}{2k + 1} b_k^2. \tag{4.3}$$

Using the result [13, p.58, Theorem 2.1.6], we have

$$b_k = \frac{2^k k!}{(2k)!} u^{(k+1)}(\eta_k), \quad \eta_k \in (-1, 1). \tag{4.4}$$

Note that $(2^k k!)/(2k)! = 1/(2k - 1)!!$. Applying the regularity assumption $|u^{(k)}(x)| \leq cM^k$, we derive

$$\begin{aligned} & \|u' - \tilde{u}'_p\|_{L_2[-1,1]}^2 \\ & < 2(cM^{p+1})^2 \left(\frac{1}{(2p - 1)!!(2p + 1)!!} + \frac{M^2}{(2p + 1)!!(2p + 3)!!} + \frac{M^4}{(2p + 3)!!(2p + 5)!!} + \dots \right) \\ & = \frac{2(cM^{p+1})^2}{(2p - 1)!!(2p + 1)!!} \left(1 + \frac{M^2}{(2p + 1)(2p + 3)} + \frac{M^4}{(2p + 1)(2p + 3)^2(2p + 5)} + \dots \right) \\ & < \frac{4(cM^{p+1})^2}{(2p - 1)!!(2p + 1)!!}, \end{aligned} \tag{4.5}$$

when $(2p + 1)(2p + 3) > 2M^2$. This last term can be readily estimated by Stirling type formula [1, (4.48)]

$$n! \approx \left(\frac{n}{e}\right)^n \sqrt{2\pi\left(n + \frac{1}{6}\right)}, \tag{4.6}$$

and [4]

$$(2n - 1)!! \approx \frac{\sqrt{(2n)!}}{\sqrt[4]{\pi\left(n + \frac{1}{4}\right)}}. \tag{4.7}$$

Therefore,

$$\begin{aligned} (2p - 1)!!(2p + 1)!! &\approx \frac{\sqrt{(2p)!}}{\sqrt[4]{\pi(p + 0.25)}} \frac{\sqrt{(2p + 2)!}}{\sqrt[4]{\pi(p + 1.25)}} \\ &\approx \frac{(2p)!2p}{\sqrt{\pi p}} \approx \left(\frac{2p}{e}\right)^{2p} 2\sqrt{2p}, \end{aligned}$$

which, combined with (4.5), leads to (4.2) with $C = \sqrt[4]{2c}$. □

Now we are ready to prove our main theorem.

Theorem 4.2. *Let $(\lambda, u) \in R^+ \times S$ be an eigen-pair of problem (2.2), where λ is a simple eigenvalue. Let $\lambda(p)$ be its approximation in the sense of (2.4) or (2.5). Then*

$$\lambda(p) - \lambda \leq Cp \left(\frac{e\sqrt{\lambda}}{4p}\right)^{2p+2}, \tag{4.8}$$

where C is independent of p and M .

Proof. Recall that $c(x)$ is constants on $(0, 1/2)$ and $(1/2, 1)$, we separate

$$\|u' - u'_p\|^2 = \|u' - u'_p\|_{L^2(0,1/2)}^2 + \|u' - u'_p\|_{L^2(1/2,1)}^2. \tag{4.9}$$

Recall that u'_p is the piecewise Legendre expansion of u' . The estimate for the first term is as follows,

$$\|u' - u'_p\|_{L^2(0,1/2)}^2 = 4\|\hat{u} - \hat{u}'_p\|_{L^2(-1,1)}^2, \tag{4.10}$$

where $\hat{u}(\xi) = u((1 + \xi)/4)$. Now we apply Lemma 4.1 to have

$$\|\hat{u} - \hat{u}'_p\|_{L^2(-1,1)} \leq C\sqrt{p} \left(\frac{e\hat{M}}{2p}\right)^{p+1} = C\sqrt{p} \left(\frac{e\sqrt{\lambda}}{4p}\right)^{p+1}.$$

Note that

$$\hat{M} = \frac{M}{4} = \frac{\sqrt{\lambda}}{2},$$

where M comes from the condition

$$\max_{x \in [0,1]} |u^k(x)| \leq cM^k.$$

In our situation, u contains terms like $\sin 2\sqrt{\lambda}x, \cos 2\sqrt{\lambda}x, \dots$ $M = 2\sqrt{\lambda}$, and $M = 4\hat{M}$ from

$$\frac{du}{dx}(x) = 4\frac{d\hat{u}}{d\xi}(\xi).$$

The second term in (4.9) can be estimated similarly, and therefore, we have

$$\|u' - u'_p\| \leq C\sqrt{p} \left(\frac{e\sqrt{\lambda}}{4p} \right)^{p+1}.$$

The error bound (4.8) follows from (4.1).

Theorem 4.3. *Let $(\lambda, u) \in R^+ \times S$ be an eigen-pair of problem (2.1), where λ is a simple eigenvalue. Let $\lambda(p)$ be its approximation in the sense of (2.4) or (2.5). Then*

$$\lambda(p) - \lambda \leq Cp \left(\frac{e\pi\sqrt{\lambda}}{2p} \right)^{2p+2}. \tag{4.11}$$

Proof. The proof is the same by the scaling between $(0, 1)$ and $(-\pi, \pi)$.

Remark. The error bounds in Theorems 4.2 and Theorem 4.3 are super-geometric type $\mathcal{O}(e^{-(2p+1)(\log p - \gamma)})$ with $\gamma = \ln(e\sqrt{\lambda}/4)$ and $\gamma = \ln(e\pi\sqrt{\lambda}/2)$, respectively. We shall demonstrate in the next section, by numerical tests, that our error bounds are sharp.

Our estimates also indicate that we need higher p for larger λ to realize the convergence. This is consistent with our numerical experiences.

5. Numerical Tests

In this section, we implement the numerical scheme described in Section 3 to solve (1.1) with $\omega = 2$ for the first 14 eigenvalues. We observe convergence for reasonably smaller p . Actually, the error goes to the machine ϵ for $p \leq 10$ for the first few eigenvalues. To verify our error bounds, we plot the ratio

$$(\lambda(p) - \lambda)/(p(0.5e\pi\sqrt{\lambda}/p)^{2p+2} \tag{5.1}$$

with some different λ s. Here is a list of the square roots of eigenvalues ($\sqrt{\lambda}$ by increasing order):

$$\begin{aligned} & \frac{1}{\pi} \arccos\left(-\frac{1}{3}\right), \quad \frac{1}{\pi} \arccos\left(-\frac{2}{3}\right), \quad \frac{1}{\pi} \arccos\frac{2}{3} + 1, \quad \frac{1}{\pi} \arccos\frac{1}{3} + 1, \quad 2, \\ & \frac{1}{\pi} \arccos\left(-\frac{1}{3}\right) + 2, \quad \frac{1}{\pi} \arccos\left(-\frac{2}{3}\right) + 2, \quad \frac{1}{\pi} \arccos\frac{2}{3} + 3, \quad \frac{1}{\pi} \arccos\frac{1}{3} + 3, \quad 4, \\ & \frac{1}{\pi} \arccos\left(-\frac{1}{3}\right) + 4, \quad \frac{1}{\pi} \arccos\left(-\frac{2}{3}\right) + 4, \quad \frac{1}{\pi} \arccos\frac{2}{3} + 5, \quad \frac{1}{\pi} \arccos\frac{1}{3} + 5, \quad 6, \dots \end{aligned}$$

Figures 1-4 demonstrate the ratio (5.1) associated with $\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_7, \lambda_9, \lambda_{14}$, respectively. We plot the ratio with different range of p . Since for a larger eigenvalue, we need relatively higher p to get into the asymptotic range. On the other hand, when p getting bigger and the error approaching the machine ϵ , the round-off error kicks in. So we can only observe the ratio in a small range of p . Nevertheless, it is sufficient to make our point clear. We see that the ratio (5.1) maintains in a reasonable range for different eigenvalues.

6. A Collocation Method for the Smooth Case

As a special case when $c(x)$ is sufficiently smooth, say a constant, we may use only one element. Without loss of generality, let us consider

$$-u'' = \lambda u \quad \text{in } (-1, 1) \quad u(-1) = 0 = u(1). \tag{6.1}$$




Fig. 5.1. Ratio of the computed errors over the estimated errors for λ_1 and λ_2

In this case, we are seeking an eigen-pair $(\lambda(p), w_p)$ with

$$w_p(\xi) = \sum_{j=2}^p w^j \hat{\phi}_j(\xi)$$

to satisfy

$$(w'_p, \hat{\phi}'_k) = \lambda(p)(w_p, \hat{\phi}_k), \quad k = 2, \dots, p. \tag{6.2}$$

Again, this result in an identity matrix on the left and a 5-diagonal matrix on the right. Based on the analysis in Section 4, we have in this case

$$\lambda(p) - \lambda \leq Cp \left(\frac{e\sqrt{\lambda}}{2p} \right)^{2p+2}. \tag{6.3}$$

Let us consider a spectral collocation method

$$-w''_p(x_j) = \lambda(p)w_p(x_j), \quad j = 1, 2, \dots, p - 1, \tag{6.4}$$

where x_j s are zeros of L'_p and L_p is the Legendre polynomial of degree p on $[-1, 1]$.

Theorem 6.1. *For the model problem (6.1), the spectral collocation method (6.4) is equivalent to replacing all integrations in (6.2) by the $p + 1$ -point Gauss-Lobatto quadrature. Furthermore, all numerical integrations are exact except the last term with*

$$(\hat{\phi}_p, \hat{\phi}_p)_* = \sum_{j=0}^p \hat{\phi}_p^2(x_j)w_j = (\hat{\phi}_p, \hat{\phi}_p) \frac{3(2p^2 - p - 1)}{2(2p^2 - p)}, \tag{6.5}$$

where w_j s are weights of the Gauss-Lobatto quadrature.

Proof. We multiply both sides of (6.4) by $\hat{\phi}_k(x_j)w_j$ and sum up

$$-\sum_{j=0}^p w''_p(x_j)\hat{\phi}_k(x_j)w_j = \lambda(p) \sum_{j=0}^p w_p(x_j)\hat{\phi}_k(x_j)w_j \tag{6.6}$$

Since the $p + 1$ -point Gauss-Lobatto quadrature rule is exact for polynomials of degree up to $2p - 1$, then we have

$$-\sum_{j=0}^p w''_p(x_j)\hat{\phi}_k(x_j)w_j = -(w''_p, \hat{\phi}_k) = (w'_p, \hat{\phi}'_k), \quad k = 2, \dots, p;$$

$$\sum_{j=0}^p w_p(x_j)\hat{\phi}_k(x_j)w_j = (w_p, \hat{\phi}_k), \quad k = 2, \dots, p - 1.$$




Fig. 5.2. Ratio of the computed errors over the estimated errors for λ_3 and λ_4




Fig. 5.3. Ratio of the computed errors over the estimated errors for λ_5 and λ_7




Fig. 5.4. Ratio of the computed errors over the estimated errors for λ_9 and λ_{14}

We see that the collocation method (6.4) is almost identical to the spectral method (6.2) except one term

$$(w_p, \hat{\phi}_p) \neq \sum_{j=0}^p w_p(x_j) \hat{\phi}_p(x_j) w_j = (w_p, \hat{\phi}_p)_*$$

and their difference is

$$(w_p, \hat{\phi}_p) - (w_p, \hat{\phi}_p)_* = w^p [(\hat{\phi}_p, \hat{\phi}_p) - \hat{\phi}_p, \hat{\phi}_p_*].$$

Using the fact $(L_p, L_p)_* = 2/p$, a direct calculation yields,

$$\begin{aligned} (\hat{\phi}_p, \hat{\phi}_p) &= \frac{1}{2(2p-1)}(L_p - L_{p-2}, L_p - L_{p-2}) \\ &= \frac{1}{2(2p-1)}((L_p, L_p) + (L_{p-2}, L_{p-2})) \\ &= \frac{1}{2(2p-1)}\left(\frac{2}{2p+1} + \frac{2}{2p-3}\right), \end{aligned} \tag{6.7}$$

and

$$\begin{aligned} (\hat{\phi}_p, \hat{\phi}_p)_* &= \frac{1}{2(2p-1)}((L_p, L_p)_* + (L_{p-2}, L_{p-2})) \\ &= \frac{1}{2(2p-1)}\left(\frac{2}{p} + \frac{2}{2p-3}\right). \end{aligned} \tag{6.8}$$

Therefore,

$$\frac{(\hat{\phi}_p, \hat{\phi}_p)_*}{(\hat{\phi}_p, \hat{\phi}_p)} = \frac{3(2p+1)(p-1)}{2p(2p-1)},$$

and (6.5) follows.

We see that the $p + 1$ -points Gauss-Lobatto quadrature has a 50% over-shoot asymptotically in calculating $(\hat{\phi}_p, \hat{\phi}_p)$. Nevertheless, the last coefficient w^p decays fast in general and the collocation method (6.4) is asymptotically equivalent to the spectral method (6.2). As a consequence, the spectral collocation method (6.4) also enjoys the super-geometric convergence rate (6.3).

Remark. It is feasible that a parallel result may be developed for the Chebyshev spectral/collocation methods. It is also feasible that the error bounds in [11] may be improved to the similar super-geometric rate as in this paper.

7. Appendix. The property of stiffness matrix

Since $(\hat{\phi}'_i, \hat{\phi}'_j) = \delta_{ij}$, it is straightforward to verify that

$$(\phi'_i, \phi'_j) = 4\delta_{ij} = (\psi'_i, \psi'_j), \quad (N', \phi'_j) = 0 = (N', \psi'_j), \quad (N', N') = 4.$$

Furthermore, observe that

$$\begin{aligned} 4(\phi_i, \phi_j) &= (\hat{\phi}_i, \hat{\phi}_j) = 4(\psi_i, \psi_j), \\ (\phi_{k+1}, N) &= 0 = (\psi_{k+1}, N) \quad k > 2, \\ (N, N) &= \frac{1}{4} \int_{-1}^1 \left(\frac{1+\xi}{2}\right)^2 + \left(\frac{1-\xi}{2}\right)^2 = \frac{1}{16} \left(4 + \frac{4}{3}\right) = \frac{1}{3}, \\ (\phi_2, N) &= \frac{1}{4}(\hat{\phi}_2, \hat{N}_2) = \frac{1}{4\sqrt{6}}(L_2 - L_0, \hat{N}_2) = \frac{-1}{4\sqrt{6}} \int_{-1}^1 \frac{1+\xi}{2} = \frac{-1}{4\sqrt{6}}, \\ (\phi_3, N) &= \frac{1}{4}(\hat{\phi}_3, \hat{N}_2) = \frac{1}{4\sqrt{10}}(L_3 - L_1, \hat{N}_2) = \frac{-1}{4\sqrt{10}} \int_{-1}^1 \xi \frac{1+\xi}{2} = \frac{-1}{12\sqrt{10}}, \\ (\psi_2, N) &= \frac{1}{4}(\hat{\phi}_2, \hat{N}_1) = \frac{1}{4\sqrt{6}}(L_2 - L_0, \hat{N}_1) = \frac{-1}{4\sqrt{6}} \int_{-1}^1 \frac{1-\xi}{2} \frac{-1}{4\sqrt{6}}, \\ (\psi_3, N) &= \frac{1}{4}(\hat{\phi}_3, \hat{N}_1) = \frac{1}{4\sqrt{10}}(L_3 - L_1, \hat{N}_1) = \frac{-1}{4\sqrt{10}} \int_{-1}^1 \xi \frac{1-\xi}{2} = \frac{1}{12\sqrt{10}}. \end{aligned}$$

The special arrangement of ordering has advantage of symmetric structure in the resulting matrix. It is straightforward to verify that the weak formulation

$$(u'_p, v') = \lambda(cu_p, v), \quad \forall v \in S_p$$

yields a diagonal stiffness matrix $4I$ and a 5-diagonal mass matrix $\frac{\lambda}{4}A$, where the upper half of A is of the form

$$\left(\begin{array}{ccccccccc} \frac{2}{(2p+1)(2p-3)} & 0 & \frac{-(2p-3)^{-1}}{(2p-1)(2p-5)} & & & & & & \\ 0 & \frac{2}{(2p-1)(2p-5)} & 0 & & & & & & \\ \frac{-(2p-3)^{-1}}{(2p-1)(2p-5)} & 0 & \frac{2}{(2p-3)(2p-7)} & \ddots & & & & & \\ & \ddots & \ddots & \ddots & \ddots & & & & \\ & & & & 0 & \frac{2}{45} & 0 & \frac{-1}{5\sqrt{21}} & \\ & & & \ddots & 0 & \frac{2}{21} & 0 & \frac{1}{3\sqrt{10}} & \\ & & & & -\frac{1}{5\sqrt{21}} & 0 & \frac{2}{5} & -\frac{1}{\sqrt{6}} & \\ & & & & & \frac{1}{3\sqrt{10}} & -\frac{1}{\sqrt{6}} & \frac{2}{3} & \end{array} \right),$$

and the lower half of A is obtained by taking the above expression up-side-down and multiplied by ω^2 . The bottom right entry $2/3$ of the upper part and the top left entry $2\omega^2/3$ of the lower part is the only overlap of the two parts.

Acknowledgement. The authors would like to thank an anonymous referee for the constructive comments (especially the hint to simplify the proof of Theorem 6.1) that significantly improves the presentation of this article. The second authors work was supported in part by the Programme for New Century Excellent Talents in University (NCET-06-0712), the National Natural Science Foundation of China (NSFC 10871066, 10571053), and Key Laboratory of High Performance Computing and Stochastic Information Processing, Ministry of Education of China, Hunan Normal University The third authors work was supported in part by the US National Science Foundation grant DMS-0612908.

References

- [1] J. Arndt and C. Haenel, π - Unleashed, Springer, Berlin, 2001.
- [2] I. Babuška and J. Osborn, Eigenvalue Problems, in: Handbook of Numerical Analysis, Vol.II: Finite Element Methods, Eds. P.G. Ciarlet and J.L. Lions, Elsevier Science Publishers B.V. (North-Holland), 1991.
- [3] I. Babuška and M. Suri, The p and $h - p$ versions of the finite element method, basic principles and properties, *SIAM Rev.*, **36** (1994), 578-632.
- [4] F.L. Bauer, Decrypted Secrets, Methods and Maxims of Cryptology, 2nd, revised, and extended edition, Springer-Verlag, Heidelberg, 2000.
- [5] C. Bernardi and Y. Maday, Spectral Methods, In Handbook of Numerical Analysis, Vol. V, P.G. Ciarlet and J.-L. Lions eds., North-Holland (1997), 209-485.
- [6] J.P. Boyd, Chebyshev and Fourier Spectral Methods, 2nd edition, Dover, New York, 2001.
- [7] C. Canuto, M.Y. Hussaini, A. Quarteroni, and T.A. Zang, Spectral Methods: Fundamentals in Single Domains, Springer Series: Scientific Computation, Berlin, 2006.

- [8] F. Chatelin, Spectral Approximations of Linear Operators, Academic Press, New York, 1983.
- [9] P.J. Davis and P. Rabinowitz, Methods of Numerical Integration, 2nd Ed., Academic Press, Boston, 1984.
- [10] D. Gottlieb and T.A. Orszag, Numerical Analysis of Spectral Methods: Theory and Applications, SIAM, Philadelphia, 1977.
- [11] M.S. Min and D. Gottlieb, Domain decomposition spectral approximations for an eigenvalue problem with a piecewise constant coefficient, *SIAM J. Numer. Anal.*, **43-2** (2005), 502-520.
- [12] A.T. Patera, A spectral element method for fluid dynamics: laminar flow in channel expansion, *J. Comput. Phys.*, **54** (1984), 468-488.
- [13] G.M. Phillips, Interpolation and Approximation by Polynomials, Springer, New York, 2003.
- [14] G. Sansone, Orthogonal Functions, Dover, New York, 1991.
- [15] J. Shen and T. Tang, Spectral and High-Order Methods with Applications, Science Press, Beijing, 2006.
- [16] L.N. Trefethen, Spectral Methods in Matlab, SIAM, Philadelphia, 2000.
- [17] Z. Zhang, Superconvergence of Spectral collocation and p-version methods in one dimensional problems, *Math. Comput.*, **74** (2005), 1621-1636.
- [18] Z. Zhang, Superconvergence of a Chebyshev spectral collocation method, *J. Sci. Comput.*, **34** (2008) 237-246.